

2

AD-A229 901

OFFICE OF NAVAL RESEARCH

Grant N00014-90-J-1193

TECHNICAL REPORT No. 32

Transient Dynamics in Excitonic Optical Bistability in Polymers

by

Xinfu Xia, Xiao-shen Li, D. L. Lin and Thomas F. George

Prepared for publication

in

Physical Review B

Departments of Chemistry and Physics  
State University of New York at Buffalo  
Buffalo, New York 14260

December 1990

DTIC  
ELECTE  
DEC 10 1990  
S E D

Reproduction in whole or in part is permitted for any purpose of the  
United States Government.

This document has been approved for public release and sale;  
its distribution is unlimited.

90 12 7 076

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFALLO/DC/90/TR-32			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Grant N00014-90-J-1193	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.		PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Transient Dynamics in Excitonic Optical Bistability in Polymers					
12. PERSONAL AUTHOR(S) Xinfu Xia, Xiao-shen Li, D. L. Lin and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day)	
				15. PAGE COUNT 18	
16. SUPPLEMENTARY NOTATION Prepared for publication in <i>Physical Review B</i>					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	* POLYMERS, ELECTRON-PHONON COUPLING		
			OPTICAL BISTABILITY, TRANSIENT DYNAMICS		
			VIRTUAL EXCITON RABI OSCILLATIONS		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Transient behavior in optical bistability mediated by phonon mode and virtual exciton coupling is studied for an electron-phonon coupling system such as polydiacetylene-toluene-sulfonate. Numerical calculations are used to analyze the evolution between bistable states, and novel switching behavior and Rabi oscillations are found.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

Physical Review B, in press

**Transient dynamics in excitonic optical bistability in polymers**

Xinfu Xia,\* Xiao-shen Li, D. L. Lin and Thomas F. George\*

Department of Physics and Astronomy  
State University of New York at Buffalo  
Buffalo, New York 14260

**Abstract**

Transient behavior in optical bistability mediated by phonon mode and virtual exciton coupling is studied for an electron-phonon coupling system such as a polydiacetylene-toluene-sulfonate. Numerical calculations are used to analyze the evolution between bistable states, and novel switching behavior and Rabi oscillations are found.

\* Also with the Department of Chemistry

1990 PACS Numbers: 42.65.-k, 42.65.Pc, 36.20.-r, 78.65.Hc

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

## 1. Introduction

Since its first observation in sodium vapor in 1976,<sup>1</sup> optical bistability has become an important phenomenon in optics, and there has been much interest in the promise of optical bistability as a mean of producing fast all-optical switching devices. Therefore, good nonlinear optical materials and more efficient device structures have been sought for practical applications, and attention is being paid to reducing sizes and decreasing the switching time and operating power for such devices.

Optical bistability means that there are two stable output optical states for a given input optical state. In general, optical bistability occurs by means of the joint effects of the nonlinear optical response of the material and the external feedback typically provided by an optical resonator. However, there are also some systems exhibiting optical bistability without the need for external feedback. For example, absorption of light tuned just below the band gap of some semiconductors can lead to increasing absorption optical bistability;<sup>2</sup> reflection at an interface between a linear and nonlinear medium<sup>3</sup> or transmission through a nonlinear thin film<sup>4</sup> have been proposed to exhibit optical bistability; and mutually induced self-focusing by interacting laser beams<sup>5</sup> and light beams interacting through four-wave mixing<sup>6</sup> have been predicted and observed, respectively, to display optical bistability. Very recently, we have found surface-induced optical bistability of polymer chains near a metal surface,<sup>7</sup> accompanied by reduced vacuum field fluctuations. In

fact, certain classes of organic molecules have been found to exhibit extremely large optical nonlinearities with fast response.<sup>8</sup> Besides, organic materials offer other attractive advantages for trying to alter the molecular structure to optimize the nonlinear optical and bulk physical properties. An example is the second-order hyperpolarizability: here the dipole of the molecular structure of a polymer is created by substituting electron-donating and electron-accepting groups at either end of the conjugated length of the molecule, where a proper choice of the donor and thus freezing in the alignment can lead to a large second-order hyperpolarizability. This flexibility in structure makes organic materials well suited for fabrication and device engineering.

Among the polymers, polydiacetylene (PDA) is a good candidate for future applications because of its giant optical nonlinearity and small transmission loss. There have been reports<sup>9</sup> about the observation of optical bistability in PDA quasi-waveguides. In the past few years, there has also been much interest in the nonresonant nonlinear optical responses of PDA-toluene sulfonate (PTS) excited by laser fields tuned well below the absorption edge.<sup>10-15</sup> Phonon-mediated bleaching<sup>11</sup> and excitonic Stark shift<sup>12</sup> have been observed, and optical nutation,<sup>13</sup> dispersive optical bistability<sup>14</sup> and splitting in pump-probe spectra<sup>15</sup> have been predicted.

In this paper, we shall theoretically analyze the transient dynamics of a PTS material irradiated by a laser field. Special attention will be paid to optical bistability and how the system

switches from one state to the other. In contrast to Ref. 14 where external feedback by an optical cavity was examined, here we shall consider another kind of bistability without external feedback. While studies of optical bistability are usually focused on the behavior of the steady-state output field intensity as a function of input intensity,<sup>2</sup> here we shall study the transient behavior of the system, through which we can extract knowledge about how the output intensity grows from zero and how fast the system evolves from one state to another,<sup>16</sup> and about possible unstable states and optical nutation. Our work is also close to experimental situations, where measurements are usually made in the transient time regime.

In Sec. II, we shall describe our model, where we treat excitons and phonons as damped oscillators and use Dekker's quantization procedure for dissipative systems<sup>17</sup> to obtain the equations of motion for the excitons and the phonons. These equations will be solved numerically for the scattered field intensity by virtual excitons, and in Sec. III we shall display the numerical results and provide a discussion and some physical conclusions.

## II. Model

It is known that there are elementary excitations of excitons and phonons in PTS<sup>18</sup> and only a few phonon modes coupled very strongly to the excitons. We consider a light beam with its frequency near the exciton resonance falling on a PTS material, and

thus the excitons interact with both the light and the lattice vibrations. The light can excite the system to create excitons and, inversely, the electrons and holes composing excitons can recombine to create radiation. However, because of the much lower frequencies of phonon modes, the number of excitons remains unchanged in their coupling with phonon modes, namely, the phonons can only play the role of mediation.<sup>11-14</sup>

It is known that both the excitons and the phonons have finite lifetimes. We may model them here as damped oscillators. For the PTS system exposed to an external electric field with frequency  $\omega$  and amplitude  $E$ , we can write a nonHermitian Hamiltonian for this system<sup>7,15</sup> as

$$H = (\omega_x - i\gamma_x) a^\dagger a + \sum_i (\omega_i - i\gamma_i) b_i^\dagger b_i + \sum_i \lambda_i a^\dagger a (b_i^\dagger + b_i) - (\mu_x a^\dagger E e^{-i\omega t} + \mu_x a E^* e^{i\omega t}) \quad (1)$$

where  $a^\dagger$  ( $a$ ) and  $b_i^\dagger$  ( $b_i$ ) stand for the creation (annihilation) operators for the exciton and  $i$ -th phonon modes with corresponding frequencies  $\omega_x$  and  $\omega_i$ , respectively,  $\gamma_x$  and  $\gamma_i$  are the damping rates for the exciton and  $i$ -th phonon mode, respectively,  $\lambda_i$  is the coupling constant for the interaction between the exciton and  $i$ -th phonon mode, and  $\mu_x$  is the component of the dipole matrix element for the exciton in the  $\vec{E}$ -field direction. As in Refs. 11-15, we neglect the momentum dependence of the exciton. With this

nonHermitian Hamiltonian characterizing energy dissipation, we can use  $a^\dagger$ ,  $a$ ,  $b_i^\dagger$  and  $b_i$  as canonical variables for this dissipative PTS system and extend Dekker's quantization procedure<sup>17</sup> for damped oscillators to write down the quantum Liouville equation for the density operator  $\rho$ :

$$\begin{aligned} \dot{\rho} = & -i[a^\dagger, [a, H]\rho] + i[\rho[H^\dagger, a^\dagger], a] - i \sum_i [b_i^\dagger, [b_i, H]\rho] \\ & + i \sum_i [\rho[H^\dagger, b_i^\dagger], b_i] \end{aligned} \quad (2)$$

In this paper, we are interested neither in any quantity that is sensitive to the quantum number counting, nor in any thermal noise present in the driving field so that quantum fluctuations can generally be neglected in our consideration.<sup>2</sup> Thus, instead of the operators, we deal with their mean values<sup>2,14</sup>  $\alpha = \langle a \rangle$  and  $\beta_i = \langle b_i \rangle$ . Such replacement simply means that we are taking the semiclassical approximation.<sup>19</sup> It has already been shown<sup>14</sup> that for excitonic optical bistability in the PTS system, there is no qualitative change in physics if we only consider a single phonon mode coupled most strongly to the excitons. Therefore, for simplicity we consider here only one phonon mode and thus delete the subscript  $i$  on  $\beta$ ,  $\omega$ ,  $\gamma$  and  $\lambda$  in the following discussion. With the equation of motion for the density operator (2), we obtain the following equations of motion for the mean values of the dynamical variables in question:



$$\dot{\alpha} = -(i\omega_x + \gamma_x)\alpha - i\lambda(\beta + \beta^*)\alpha + i\mu_x^* E e^{-i\omega_L t} , \quad (3a)$$

$$\dot{\beta} = -(i\omega + \gamma)\beta - i\lambda|\alpha|^2 . \quad (3b)$$

For convenience, in what follows we use the rotating frame.

Namely, we replace  $\alpha$  in Eqs. (3) by  $\alpha e^{-i\omega_L t}$  to obtain

$$\dot{\alpha} = -(i\Delta + \gamma_x)\alpha - i\lambda(\beta + \beta^*)\alpha + i\Omega , \quad (4a)$$

$$\dot{\beta} = -(i\omega + \gamma)\beta - i\lambda|\alpha|^2 , \quad (4b)$$

where we have defined the detuning  $\Delta = \omega_x - \omega_L$  and the Rabi frequency  $\Omega = \mu_x^* E$ .

When the external laser field with its frequency near the exciton resonance falls on the PTS material, the radiation field is scattered by the excitons. In the scattering region where the incident external field vanishes, the relation between the positive-frequency part of the scattered field and the excitonic dipole moment can be written as<sup>20</sup>

$$E_{sc}^{(+)}(\vec{r}) = \psi(\vec{r})a(t - \frac{r}{c}) , \quad (5)$$

where  $\psi(\vec{r}) = -\frac{\omega_x^2}{4\pi c^2 r^3} (\vec{\mu}_x \times \vec{r}) \times \vec{r}$ , and  $\vec{\mu}_x a$  is the excitonic dipole moment. Hence, the mean value of the dimensionless scattered field intensity is given by

$$I_{sc}(t) = \langle |E_{sc}^{(+)}(\vec{r}, t)|^2 \rangle / \psi^2(\vec{r}) = |\alpha|^2 . \quad (6)$$

In the steady state, it is easy to find, by setting the terms on the left-hand side of Eqs. (4) equal to zero, that this scattered intensity,  $I_{sc}(t \rightarrow \infty)$ , is determined by the cubic equation

$$\lambda_p^2 I_{sc}^3 - 2\Delta \lambda_p I_{sc}^2 + (\Delta^2 + \gamma_x^2) I_{sc} - I_{in} = 0 , \quad (7)$$

where we have defined the incident field intensity  $I_{in} = |\Omega|^2$ , and  $\lambda_p = 2\lambda^2\omega/(\omega^2 + \gamma^2)$ . However, in the transient regime, we have to solve the nonlinear Eqs. (4) numerically.

### III. Results and discussion

In this paper we use the following parameters for PTS:<sup>11,14</sup>  
 $\gamma_x = 0.05$  eV,  $\gamma = 0.002$  eV,  $\lambda = 0.1$  eV and  $\omega = 0.258$  eV. For the first step, it is helpful to solve for the steady-state solution using Eq. (7). Through analyzing this equation, we have found that

only when the incident field is tuned sufficiently below the exciton resonance, say,  $\omega_x - \omega_L > 0.4\omega$ , does there exist optical bistability, indicating that the virtual exciton is essential for this bistability. We have also found that the phonon-exciton coupling is indispensable to this bistability. This is similar to the case where PTS is placed in an optical cavity.<sup>14</sup>

Figure 1(a) is an example showing the scattered field intensity plotted vs the incident field intensity when the system is in steady state and  $\Delta = 0.5\omega$ . Optical bistability is clearly shown in this figure. Generally speaking, optical bistability occurs only when the system exhibits nonlinear responses to the external field and a feedback of the field or other agent exists. In semiconductors, it is well known that<sup>10,11</sup> virtual excitons play almost the same role as do real excitons in response to external fields. In the present case, as the incident field intensity changes, the dynamics of virtual excitons changes accordingly, leading to a subsequent change in the phonon mode. As a consequence, the exciton-phonon coupling changes the properties of the virtual excitons, resulting in a change in the scattered light. This internal process of "feedback" provided by the phonon mode permits possible multivalued solutions to the output intensity of the nonlinear optical scattering problem. This means that within a certain range, there can be three distinct scattered intensities for each incident intensity. We have found, through numerical simulation, that the middle solution is unstable, and hence only the upper and lower solutions can be reached. As observed in Fig.

1(a), when  $I_{in}$  increases continuously from zero,  $I_{sc}$  increases continuously until it reaches point b where it becomes unstable. Further increase of  $I_{in}$  leads to an abrupt jump of  $I_{sc}$  to the upper branch. Similarly, when  $I_{in}$  decreases continuously,  $I_{sc}$  jumps to the lower branch at point a where the solution is unstable.

Through numerical calculations of Eqs. (4) and (6), we have studied the time evolution of the scattered intensity after a sudden intensity change or sudden switching on of the incident field. Figures 1(b), 1(c) and 2(a)-(c) show some interesting results. Figure 1(b) describes the situation that when  $I_{sc}$  reaches steady state (point a) with  $I_{in} = 1.58\gamma_x^2$ , then we suddenly increase  $I_{in}$  from  $1.58\gamma_x^2$  to  $2.29\gamma_x^2$ . How  $I_{sc}$  evolves from the lower state (point a) to the upper state (point c) is clearly shown. We see that the switching time is approximately one order of magnitude longer than the lifetime of the virtual exciton but the same order of magnitude as the lifetime of the phonon mode. Comparing Fig. 1(a) with 1(b), one can find that in the course of  $I_{sc}$  going from the lower state to the upper one (from point a to c), it goes along the line of the lower branch, and then gradually goes upward, and finally, after oscillating a bit, reaches point c. Figure 1(c) describes the reverse case of Fig. 1(b). By comparing Figs. 1(b) and 1(c), one can see that  $I_{sc}$  goes along a different path from point c to a. By comparing Figs. 1(a) and 1(c), one can see that  $I_{sc}$  goes along the upper branch until it reaches point d, then it goes gradually down to point a. Therefore, interestingly enough, through our study in the transient regime, we can obtain detailed

information about not only the evolution of  $I_{sc}$  between bistable states, but also how it completes a hysteresis loop.

In addition to the switching behavior of the scattered field intensity  $I_{sc}$ , we have also studied how  $I_{sc}$  builds up from zero through a switching-on by an incident field at initial time  $t = 0$ . Some numerical results calculated from Eqs. (4) and (6) are shown in Fig. 2, in which we have chosen three different input intensities in the region that bistability occurs. For fixed detuning  $\Delta$  there is a corresponding threshold, and smaller  $\Delta$  leads to lower threshold as long as bistability occurs. If  $I_{in}$  lies below the threshold,  $I_{sc}$  eventually evolves to reach the lower state. Otherwise,  $I_{sc}$  eventually reaches the upper state. In Fig. 2 we choose  $\Delta = 0.5\omega$ , and the subsequent threshold for  $I_{in}$  is found to be approximately  $2.21\gamma_x^2$ . By comparing Fig. 1(a) with Fig. 2, we see that in Figs. 2(a) and (b) where the  $I_{in}$ 's are smaller than this threshold value,  $I_{sc}$  goes into the lower branch, while in Fig. 2(c) where  $I_{in}$  is beyond the threshold,  $I_{sc}$  goes into the upper branch. From Fig. 2 we can also see Rabi oscillations. However, these oscillations are modulated, i.e., the centers of oscillation change with respect to time. When  $I_{in}$  is below the threshold,  $I_{sc}$  first goes up then goes down to reach the lower state [see Figs. 2(a) and (b)]. By contrast, when  $I_{in}$  is above the threshold,  $I_{sc}$  first reaches the lower state then goes up to reach the upper state [see Fig. 2(c)]. In addition, through numerical analysis, we have found that no matter what initial condition is used,  $I_{sc}$  cannot evolve into the middle branch [see the line between points b and d in Fig.

1(a)]. So, we can conclude that the middle or third solution of  $I_{sc}$  is not stable. Meanwhile, the peaks in Figs. 2(a) and (b) show the possibility for  $I_{sc}$  to temporarily stay in the middle state, which means that the middle state can only be reached temporarily.

#### Acknowledgment

This work was supported by the Office of Naval Research.

## References

1. H. M. Gibbs, S. L. McCall and T. N. C. Venkatesan, Phys. Rev. Lett. 36, 1135 (1976).
2. H. M. Gibbs, Optical Bistability: Controlling Light with Light (Academic, New York, 1985).
3. A. E. Kaplan, JETP Lett. 24, 114 (1976).
4. W. Chen and D. L. Mills, Phys. Rev. B 38, 12814 (1988).
5. A. E. Kaplan, Opt. Lett. 6, 360 (1981).
6. D. J. Gauthier, M. S. Malcuit, A. L. Gaeta and R. W. Boyd, Phys. Rev. Lett. 64, 1721 (1990).
7. D. L. Lin, X. S. Li and T. F. George, Phys. Lett. A, submitted.
8. See, for example, D. S. Chemla and J. Zyss. Eds., Nonlinear Optical Properties of Organic Molecules and Crystals (Academic, New York, 1987).
9. B. P. Singh and P. N. Prasad, J. Opt. Soc. Am. B 5, 453 (1988); K. Sasaki, K. Fujii, T. Tomioka and T. Kinoshita, J. Opt. Soc. Am. B 5, 457 (1988).
10. B. I. Greene, J. Orenstein, R. R. Millard and L. R. Williams, Chem. Phys. Lett. 139, 381 (1987); Phys. Rev. Lett. 58, 2750 (1987).
11. B. I. Greene, J. F. Mueller, J. Orenstein, D. H. Rapkine, S. Schmitt-Rink and M. Thakur, Phys. Rev. Lett. 61, 325 (1988); B. I. Greene, J. Orenstein and S. Schmitt-Rink, Science 247, 679 (1990).

12. G. J. Blanchard, J. P. Heritage, A. C. Von Lehmen, M. K. Kelly, G. L. Baker and S. Etemad, Phys. Rev. Lett. 63, 887 (1989).
13. X. S. Li, D. L. Lin, T. F. George and X. Sun, Phys. Rev. B 40, 11728 (1989).
14. X. S. Li, D. L. Lin, T. F. George and X. Sun, Phys. Rev. B (Rapid Commun.) 41, 3280 (1990); Phys. Rev. B, in press.
15. X. Xia, X. S. Li, D. L. Lin and T. F. George, Phys. Rev. B (Rapid Commun.), in press.
16. See, for example, V. Benza and L. A. Lugiato, Lett. Nuovo Cimento, 26, 405 (1979).
17. H. Dekker, Physica A 95, 311 (1979).
18. D. N. Batchelder, in Polydiacetylenes, edited by D. Bloor and R. R. Chance (Martins Nijhoff, Dordrecht, The Netherlands, 1985), p. 187 ff.
19. M. Reid, K. J. McNeil and D. F. Walls, Phys. Rev. A 24, 2029 (1981).
20. B. R. Mollow, Phys. Rev. 188, 1969 (1969).



### Figure Captions

1. (a) Scattered field intensity  $I_{sc}$  vs driving field intensity  $I_{in}$  (in the unit of  $\gamma_x^2$ ) for steady-state operation. (b) Transition of  $I_{sc}$  from the lower state, corresponding to  $I_{in} = 1.58\gamma_x^2$ , to the upper state, corresponding to  $I_{in} = 2.29\gamma_x^2$ . (c) Transition of  $I_{sc}$  from the upper state corresponding to  $I_{in} = 2.29\gamma_x^2$ , to the lower state, corresponding to  $I_{in} = 1.58\gamma_x^2$ .
  
2. Time (t) evolution of the scattered field intensity  $I_{sc}$  for different driving intensities, where (a)  $I_{in} = 1.93\gamma_x^2$ , (b)  $I_{in} = 2.20\gamma_x^2$  and (c)  $I_{in} = 2.21\gamma_x^2$ .

Fig. 1







Fig. 2



FY90\* Abstracts Distribution List for Solid State & Surface Chemistry

Professor John Baldeschwieler  
Department of Chemistry  
California Inst. of Technology  
Pasadena, CA 91125

Professor John Eyler  
Department of Chemistry  
University of Florida  
Gainesville, FL 32611

Dr. Sylvia Johnson  
SRI International  
333 Ravenswood Avenue  
Menlo Park, CA 94025

Professor Paul Barbara  
Department of Chemistry  
University of Minnesota  
Minneapolis, MN 55455-0431

Professor James Garvey  
Department of Chemistry  
State University of New York  
Buffalo, NY 14214

Dr. Zakya Kafafi  
Code 6551  
Naval Research Laboratory  
Washington, DC 20375-5000

Dr. Duncan Brown  
Advanced Technology Materials  
520-D Danury Rd.  
New Milford, CT 06776

Professor Steven George  
Department of Chemistry  
Stanford University  
Stanford, CA 94305

Professor Larry Kesmodel  
Department of Physics  
Indiana University  
Bloomington, IN 47403

Professor Stanley Bruckenstein  
Department of Chemistry  
State University of New York  
Buffalo, NY 14214

Professor Tom George  
Dept. of Chemistry and Physics  
State University of New York  
Buffalo, NY 14260

Professor Max Lagally  
Dept. Metal. & Min. Engineeri  
University of Wisconsin  
Madison, WI 53706

Professor Carolyn Cassady  
Department of Chemistry  
Miami University  
Oxford, OH 45056

Dr. Robert Hamers  
IBM T.J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, N Y 10598

Dr. Stephen Lieberman  
Code 522  
Naval Ocean Systems Center  
San Diego, CA 92152

Professor R.P.H. Chang  
Dept. Matls. Sci. & Engineering  
Northwestern University  
Evanston, IL 60208

Professor Charles Harris  
Department of Chemistry  
University of California  
Berkeley, CA 94720

Professor M.C. Lin  
Department of Chemistry  
Emory University  
Atlanta, GA 30322

Professor Frank DiSalvo  
Department of Chemistry  
Cornell University  
Ithaca, NY 14853

Professor John Hemminger  
Department of Chemistry  
University of California  
Irvine, CA 92717

Professor Fred McLafferty  
Department of Chemistry  
Cornell University  
Ithaca, NY 14853-1301

Dr. James Duncan  
Federal Systems Division  
Eastman Kodak Company  
Rochester, NY 14650-2156

Professor Leonard Interrante  
Department of Chemistry  
Rensselaer Polytechnic Institute  
Troy, NY 12181

Professor Horia Metiu  
Department of Chemistry  
University of California  
Santa Barbara, CA 93106

Professor Arthur Ellis  
Department of Chemistry  
University of Wisconsin  
Madison, WI 53706

Professor Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, NY 14853

Professor Larry Miller  
Department of Chemistry  
University of Minnesota  
Minneapolis, MN 55455-0431

Professor Mustafa El-Sayed  
Department of Chemistry  
University of California  
Los Angeles, CA 90024

Professor Eugene Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, NC 27514

Professor George Morrison  
Department of Chemistry  
Cornell University  
Ithaca, NY 14853

Professor Daniel Neumark  
Department of Chemistry  
University of California  
Berkeley, CA 94720

Professor Robert Whetten  
Department of Chemistry  
University of California  
Los Angeles, CA 90024

Professor David Ramaker  
Department of Chemistry  
George Washington University  
Washington, DC 20052

Professor R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, CA 90024

Dr. Gary Rubloff  
IBM T.J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, NY 10598

Professor Nicholas Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, PA 16902

Professor Richard Smalley  
Department of Chemistry  
Rice University  
P.O. Box 1892  
Houston, TX 77251

Professor Aaron Wold  
Department of Chemistry  
Brown University  
Providence, RI 02912

Professor Gerald Stringfellow  
Dept. of Matls. Sci.  
& Engineering  
University of Utah  
Salt Lake City, UT 84112

Professor Vicki Wysocki  
Department of Chemistry  
Virginia Commonwealth University  
Richmond, VA 23284-2006

Professor Galen Stucky  
Department of Chemistry  
University of California  
Santa Barbara, CA 93106

Professor John Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburg, PA 15260

Professor H. Tachikawa  
Department of Chemistry  
Jackson State University  
Jackson, MI 39217-0510

Professor William Unertl  
Lab. for Surface Sci.  
& Technology  
University of Maine  
Orono, ME 04469

Dr. Terrell Vanderah  
Code 3854  
Naval Weapons Center  
China Lake, CA 93555

Professor John Weaver  
Dept. of Chem. Eng. & Mat. Sci.  
University of Minnesota  
Minneapolis, MN 55455

Professor Brad Weiner  
Department of Chemistry  
University of Puerto Rico  
Rio Piedras, Puerto Rico 00931